## FastDigest Restriction Enzymes





## One buffer for 176 enzymes—it's that easy

Thermo Scientific<sup>™</sup> FastDigest<sup>™</sup> Restriction Enzymes support complete and fast digestions.

#### Why use FastDigest enzymes?

FastDigest enzymes are an advanced line of restriction enzymes that offer fast and complete digestion of DNA in a single, universal buffer.

#### Why use the colorless buffer?

We recommend using the colorless Thermo Scientific™ FastDigest™ Buffer for applications that require product analysis by fluorescence excitation (e.g., concentration measurements in UV light).

#### Why use the green buffer?

Thermo Scientific™ FastDigest™ Green Buffer allows for direct loading of the reaction mixture on gels. The green buffer contains a density reagent and two tracking dyes that do not interfere with downstream applications, including dephosphorylation, endrepair reactions, and ligation.


### Thermo Fisher SCIENTIFIC



# 176 FastDigest Restriction Enzymes universal system

- 1 buffer, 2 formats
- Complete digestion in 5–15 min
- No star activity due to short incubation times
- Direct loading on gel

### **New England BioLabs**



#### 276 restriction enzymes

#### Abbreviations used:

CS = CutSmart<sup>™</sup> buffer—universal buffer
TSQ = Time-Saver<sup>™</sup> qualified; 5–15 min digestion
HF = High-Fidelity (HF<sup>™</sup>) enzymes; engineered for reduced star activity

#### Features (Figure 1):

- The first restriction enzyme offering using a universal buffer
- Double and multiple digestions in a universal buffer for any combination of enzymes
- No sequential digestions and buffer changes
- 176 unique specificities
- Complete digestion in 5–15 minutes
- Direct loading of reaction mixture on gels

Figure 1. Simplicity of FastDigest restriction enzymes, a universal system using one buffer vs. the complicated NEB™ offering. Note, this is an illustrative image depicting the comparison of features between the two companies.

**Thermo Fisher** s c i e n t i f i c

#### **Technical details**

- 1 μL of FastDigest enzyme cleaves 1 µg of substrate DNA in 5-15 minutes in FastDigest buffer (Figure 2)
- Designed to eliminate star activity due to short incubation times
- All enzymes qualified for rapid and complete digestion of all types of DNA
- · Protocols for plasmid, genomic, and viral DNA as well as PCR products are provided


#### **Usage and applications**

Choose FastDigest enzymes for traditional molecular cloning techniques, including:

- Clone analysis
- Preparation of DNA for cloning
- Digestion of PCR products
- Restriction fragment length polymorphism (RFLP) genotyping
- Digestion of difficult-to-cleave DNA

Α


#### Plasmid DNA digested with FastDigest enzymes



- 1: undigested plasmid DNA
- 2: FastDigest Bcul
- 3: FastDigest Xbal
- 4: FastDigest Ndel
- 5: FastDigest Sall
- 6: FastDigest XmaJI
- 7: FastDigest Eco31I
- 8: FastDigest Eco52I
- 9: FastDigest BgIII

В

#### Plasmid DNA digested with NEB enzymes



- 1: undigested plasmid DNA
- 2: Spel-HF
- 3: Xbal
- 4: Ndel
- 5: Sall-HF
- 6: AvrII
- 7: Bsal
- 8: Eagl-HF
- 9: Bgll

Figure 2. Comparison of digestion efficiencies of restriction enzymes. (A) FastDigest restriction enzymes digest plasmid DNA much more efficiently compared to the (B) NEB enzymes. In this experiment, using the NEB protocol, 1 µg of plasmid DNA was digested in ~15 minutes.

#### **Ordering information**

For a complete list of the 176 FastDigest enzyme specificities, visit thermofisher.com/fastdigest

To try our top 13 enzymes, and both the colorless and green FastDigest Buffers, get our Thermo Scientific™ FastDigest™ Value Pack (Cat. No. K1991), or visit thermofisher.com/fdvaluepack

#### Buffer compatibility with downstream applications

Thermo Scientific™ DNA/RNA modifying enzymes	Activity in FastDigest Green Buffer or FastDigest Buffer (colorless)
DNA Polymerase I, E. coli	100%
Klenow Fragment	100%
Klenow Fragment, exo-	100%
T7 DNA Polymerase	100%
T4 DNA Ligase*	75–100%
FastAP™ Thermosensitive Alkaline Phosphatase	100%
T4 Polynucleotide Kinase (T4 PNK)	100%

<sup>\* 0.5</sup> mM ATP is required for T4 DNA ligase activity.

## 000 «Диаэм»

Москва ул. Магаданская, д. 7, к. 3 ■ тел./факс: (495) 745-0508 ■ sales@dia-m.ru

www.dia-m.ru

С.-Петербург +7 (812) 372-6040 spb@dia-m.ru

Казань +7(843) 210-2080 kazan@dia-m.ru

Новосибирск +7(383) 328-0048 nsk@dia-m.ru

Ростов-на-Дону +7 (863) 303-5500 rnd@dia-m.ru

Воронеж +7 (473) 232-4412 vrn@dia-m.ru

Екатеринбург +7 (912) 658-7606 ekb@dia-m.ru

Йошкар-Ола +7 (927) 880-3676 nba@dia-m.ru

Кемерово +7 (923) 158-6753 kemerovo@dia-m.ruu

Красноярск +7(923) 303-0152 krsk@dia-m.ru

**Армения** +7 (094) 01-0173 armenia@dia-m.ru

